

AMREF INTERNATIONAL UNIVERSITY SCHOOL OF MEDICAL SCIENCES DEPARTMENT OF NURSING AND MIDWIFERY SCIENCES KENYA REGISTERED COMMUNITY HEALTH NURSING END OF SEMESTER EXAMINATIONS

DNS 122: MEDICAL SURGICAL NURSING I

DATE: 17TH **A**PRIL 2024

TIME:2 Hours START: 0900 HOURS FINISH: 1100 HOURS

INSTRUCTIONS

- 1. This exam will be marked out of 70 Marks
- **2.** ALL Questions are compulsory.
- **3.** The Examination has Three Sections: Section I- Multiple Choice Questions, Section II: Short Answer Questions, Section III: Long Essay Questions
- **4.** Answer all Questions in the ANSWER BOOKLET provided
- **5.** Do Not write anything on the question paper -use the back of your booklet for rough work if need be.

SECTION A: MULTIPLE CHOICE QUESTIONS (30 MARKS)

- 1. The appropriate position for a client undergoing paracentesis is:
 - a. Prone.
 - b. Supine
 - c. Side-lying, with legs pulled up and head bent down onto the chest
 - d. Side-lying, with a pillow under the hip, and head straightened
- 2. To prevent post procedural complications associated with lumbar puncture:
 - **a.** Observe strict asepsis, ensure patient remains flat 6-8 hours after the procedure
 - b. Limit fluid intake, nurse in a bright airy room
 - c. Maintain strict asepsis, position client in semi fowlers position
 - d. Encourage ambulation within 2 hours, Limit fluid intake
- 3. The type biopsy recommended for taking a bone marrow sample is the:
 - a. Aspiration biopsy
 - b. Punch biopsy
 - c. Excisional biopsy
 - d. Brush biopsy
- **4.** Pharmacologic agents that control pain by interfering with depolarization of the nerve include:
 - a. Non steroidal Anti Inflammatory Drugs (NSAIDS)
 - **b.** Corticosteriods
 - c. Anticonvulsants
 - **d.** Opioid analgesics
- 5. The following constitutes the body's first line of defense:
 - a. Tears
 - b. T lymphocytes
 - **c.** The complement system
 - d. Neutrophils

- 6. In wound healing:
 - **a.** Healing by primary intention occurs when wound margins cannot be approximated
 - b. Chemotherapy can delay wound healing
 - c. Healing by secondary intention occurs with minimal scarring
 - d. Clean surgical wounds heal by tertiary intention
- 7. In the nursing care of a client with fluid volume excess:
 - a. Weight gain is the best indicator of fluid retention
 - b. Edema is objectively estimated by a visual assessment
 - c. The risk of impaired skin integrity is minimal
 - d. Interventions to support respiratory status are not a priority
- 8. Insensible fluid losses include:
 - a. Urine
 - b. Gastric drainage
 - c. Bleeding
 - d. Perspiration
- **9.** Nursing interventions to manage pain in a client who cannot communicate verbally includes:
 - a. Assess the nonverbal cues of pain to establish severity.
 - b. Obtain baseline behavioral indicators from family members.
 - c. Review medical records for the time ,dose and response to the last dose of treatment
 - d. Administer a bolus dose of opioid analgesics immediately
- **10.** Using the parklands formula and 3mls/kg bwt, calculate the amount of fluid to be replaced within the first 8 hours to an 80kg patient who sustains burns to the entire right arm and chest:
 - a. 4320
 - b. 2160
 - **c.** 1080
 - **d.** 8640

- 11. Hemophilia is characterized by a deficiency of :
 - a. Iron
 - b. White blood cells
 - c. Factor VIII
 - d. Vitamin K
- **12.** Nursing care for patient with a platelet count of 25,000 per microliter of blood should include:
 - a. Close monitoring for Fever, signs of infection
 - b. Transfusion with packed red blood cells
 - **c.** Monitoring for bleeding episodes
 - d. Reverse barrier nursing
- 13. Iron deficiency anaemia
 - a. Is characterized by microcytic erythrocytes
 - b. Is characterized by megaloblastic erythrocytes
 - **c.** In children is caused by tapeworm infestation
 - **d.** The gold standard of treatment is transfusion with whole blood.
- **14.** Muscle wasting in a patient with Cushing's syndrome can be reduced by administering;
 - a. High calorie diet
 - b. High protein diet
 - c. Vitamin D
 - d. Low sodium diet
- 15. Nursing interventions for a patient with hyperparathyroidism include:
 - a. Fluid restriction
 - b. Increase intake of calcium rich food
 - **c.** Encourage ambulation
 - **d.** Nursing the patient in a dim lighted room
- **16.** Pituitary gland disorder that results in reduced production of Adrenocorticotropic hormone (ACTH) may lead to:
 - a. Diabetes insipidus
 - b. Cushing's syndrome
 - c. Addison's disease
 - d. Hypogonadism

- 17. A patient with Addison's disease is likely to present with:
 - a. Hypertension
 - b. Hyperpigmentation
 - c. Hyperglycemia
 - d. Hypernatremia
- 18. A pituitary tumour secreting excess vasopressin is likely to cause:
 - **a.** Diabetes inspidus
 - b. Cretinism
 - c. Syndrome of inappropriate antidiuretic hormone secretion
 - d. Gigantism
- 19. Interventions for a patient with Hypothyroidism include:
 - a. Nursing the patient in a cool room
 - **b.** Preparing the patient for thyroidectomy
 - c. Administration of laxatives
 - d. Administering sedatives
- 20. The nutritional interventions for a patient with hyperthyroidism include:
 - a. Provide privacy during meal times
 - b. Encourage the patient to reduce the amount of food they eat
 - **c.** Minimise fluid intake as much as possible
 - **d.** A high fat diet is encouraged
- 21. Pheochromocytomas mainly secrete:
 - a. Cortisol
 - b. Adrenaline
 - c. Aldosterone
 - d. Androgens
- 22. Heart failure is:
 - **a.** Inability of the heart to fit in the mediastinum
 - b. Limited blood flow to the heart due to severe congestion
 - **c.** Inability of the heart to pump blood that meets body tissue needs
 - **d.** Dextraposition of the heart alongside myocardial hypertrophy

- **23.** To prevent Coronary Heart Disease (CHD), the following diet should be encouraged:
 - a. Plant proteins and animal fats
 - b. Soluble fiber and omega 3 fatty acids
 - c. Seasoned foods and sea foods
 - d. Triglycerides and plant proteins
- 24. The following is a danger sign for a patient with chest injury;
 - a. Systolic BP of < 120mmhg
 - b. Heart Rate of > 100 Beats / min
 - c. PaCo2 > 5kpa on room air
 - d. Reducing size of a pneumothorax
- 25. The following is a common predisposing factor to aneurysms:
 - a. Male gender
 - b. Hypertension
 - c. Repeated typhoid infections
 - d. Diet rich in nitrosamines
- 26. A tension pneumothorax is,
 - a. A complication of a simple pneumothorax
 - b. The commonest cause of myocardial infarction
 - **c.** Preferably diagnosed clinically
 - d. Preferably diagnosed through a chest xray
- 27. The most common predisposing factor for Asthma is:
 - a. Genetics
 - b. Age
 - c. Cold temperatures
 - d. Dust
- 28. The following breath sound is a cardinal sign of an acute asthmatic attack:
 - a. Crackle
 - b. Stridor
 - c. Rale
 - d. Wheeze

- 29. The most common microorganism that causes high incidence of Pneumonia is:
 - a. Staphylococcus aureus
 - b. Streptococcus Pneumoniae
 - **c.** Covid 19
 - d. Pneumocystis carinii pneumonia
- 30. The following condition mainly manifests with a barrel chest:
 - a. Emphysema
 - b. Pumonary tuberculosis
 - **c.** Bronchial asthma
 - **d.** Pharyngitis

SECTION II: SHORT ANSWER QUESTIONS (20 MARKS)

- 1. Outline five (5) factors that interfere with wound healing (5 Marks)
- 2. State four (4) non pharmacological measures that can be instituted to control pain (4 Marks)
- 3. Describe the nursing management of a patient with an Acute Vaso occlusive crisis in sickle cell disease (5 Marks)
- 4. Giving examples, describe any three (3) classifications of shock (6 Marks)

SECTION III: LONG ANSWER QUESTION (20 MARKS)

- 1. Mr. Q, 50 yrs old is admitted with left sided heart failure.
 - a. Draw and label a diagram illustrating electrical conductivity of the heart (4 Marks)
 - b. Describe the pathophysiology of heart failure (4 Marks)
 - c. Describe the management of Mr Q from admission till discharge (12 Marks)