

**091906T4HRI**

**Health Records and Information Technology Level 6**

**AITC/HRIT/03/6/A**

**Biostatistics and Data Analysis. (Medical Statistics 2)**

**July/August 2025**



**AMREF INTERNATIONAL TRAINING CENTER**

**CANDIDATE WRITTEN ASSESSMENT**

**Time: 3 HOURS**

**INSTRUCTIONS TO CANDIDATE**

1. Marks for each question are indicated in the brackets.
2. The paper consists of **TWO** sections: **A** and **B**.
3. Candidates are provided with a separate answer booklet
4. **DO NOT** write on this question paper.

**This paper consists of THREE (3) printed pages**

**Candidates should check the question paper to ascertain that all pages are printed as indicated and that no questions are missing.**

**SECTION A: (40 MARKS)***Answer all questions in this section.*

1. Define the following terms.
  - i) Biostatistics. (2 Marks)
  - ii) Level of significance. (2 Marks)
2. Differentiate the following terms.
  - i) Type I error and Type II error. (2 Marks)
  - ii) Homoscedasticity and heteroscedasticity. (2 Marks)
  - iii) Null hypothesis and alternative hypothesis. (2 Marks)
3. Using the given health-related dataset (18, 34, 76, 29, 15, 41, 46, 25, 54, 38, 20, 32, 43, 22), Construct a box-and-whisker plot (box plot) to summarize the distribution of the data and identify any potential outliers. (3 Marks)
4. Consider the following set;  $x = \{11,13,15,16,19,22,13,20\}$   
Calculate;
  - i) Arithmetic mean. (2 Marks)
  - ii) Harmonic mean. (2 Marks)
5. A researcher analyzed the association between injury severity (fatal vs. non-fatal) across two groups using a Chi-square test in R. The test results were as follows:

```

fatal=c(1601,510)
non_fatal=c(162527,412368)
status=cbind(fatal,non_fatal)
chisq=chisq.test(status)
chisq

##
## Pearson's Chi-squared test with Yates' continuity correction
##
## data:  status
## X-squared = 2336.1, df = 1, p-value < 2.2e-16

```

- i) Based on the output above, is the result statistically significant at the 5% level? Justify your answer. (2 Marks)
- ii) What does the p-value tell us about the relationship between fatal and non-fatal outcomes across the two groups? (2 Marks)
- 6. Discuss TWO properties of a student t distribution. (2 Marks)
- 7. Suppose a health facility is interested in the average recovery time (in days) for patients after a specific treatment. A random sample of 36 patient records shows a mean recovery time of 68 days. If the population standard deviation is known to be 3 days, calculate the 95% confidence interval estimate for the average recovery time of all patients who undergo this treatment. (3 Marks)
- 8. Consider the following ANOVA (Analysis of Variance) table. Copy and complete the ANOVA table. (3 Marks)

Source of Variation	Degrees of Freedom (df)	Sum of Squares (SS)	Mean Sum of Squares (MS)	f
Regression	1	.....	799.14	.....
Residual	.....	215.61	107.81	
Total	.....	.....		

- 9. Outline TWO types of statistics commonly applied within healthcare organizations to support decision-making and performance evaluation. (3 Marks)
- 10. Two samples of patient recovery times (in days) after receiving different treatment methods are taken from normally distributed populations. The first sample has a mean recovery time of 4.29 days and a variance of 6.89, based on 10 patients. The second sample has a mean of 4.18 days and a variance of 2.55, based on 12 patients. Using this data:

- i) State the null and alternative hypotheses to determine if there is a significant difference in mean recovery times between the two treatment methods. (2 Marks)
- ii) Identify the appropriate statistical test and justify your results. (2 Marks)

11. In 2024, Shianda hospital had a bed complement of 120 and an average length of stay of 6 days. It also had a daily occupancy of 105 with a bed turnover of 54 patients. This turnover is expected to increase by 12% this year (2025). Using this data, calculate the following information for Shianda;

- i) Available bed days. (2 Marks)
- ii) Occupied bed days. (2 Marks)

AMREF

**SECTION B: (60 MARKS)***Answer any THREE questions in this section*

12.

- a) In a recent public safety and health assessment conducted by the County Government of Trans Nzoia, researchers collected data to investigate the relationship between alcohol consumption and types of crimes committed. The study aimed to provide insights into whether alcohol use might influence the prevalence or nature of criminal activity within the county. The data collected included six categories of crimes and the number of individuals involved in each crime, classified according to their alcoholic status (Alcoholic or Non-Alcoholic). The summarized findings are presented in the table below:

Crime	Alcoholic	Non -Alcoholic
Arson	50	43
Rape	88	62
Violence	155	110
Theft	379	300
Coning	18	14
Fraud	63	144

As a health analyst, you are tasked with evaluating whether alcoholic status has a significant association with the type of crime committed. Your analysis will assist policymakers in designing targeted interventions.

- i) Determine whether or not Alcoholic status and Type of Crime are dependent. (8 Marks)
- ii) Provide the R code that can be used to determine whether there is a significant association between alcoholic status and type of crime in Trans Nzoia County. (2 Marks)

- b) As part of a community health survey, Amina, a community health officer, collected data on households' health awareness scores (rated from 0 to 100). The scores were grouped as follows:

Health Score Range	Number of Households (f)
50-59	3
60-69	5
70-79	9
80-89	12
90-100	8

Using the data, calculate the following measures to assist Amina in evaluating the overall health awareness of the community:

- i) Mean health awareness score. (2 Marks)
- ii) Variance. (4 Marks)
- iii) Standard deviation. (2 Marks)
- iv) Give an R code for calculating the mean, variance and standard deviation for the above data. (2 Marks)

13.

- a) The variables X and Y represent health assessment scores recorded during a preliminary screening and final evaluation, respectively, for patients enrolled in a medical outreach program by Amref Flying Doctors.

X (Preliminary Score)	Y Final score
70	77
74	94
72	88
68	80
58	71
54	76
82	88
64	80
80	90
61	69

Using the above patient data collected by Amref Flying Doctors, do the following:

- i) Draw a scatter plot to examine the relationship between preliminary and final health assessment scores. (2 Marks)
- ii) Fit a simple linear regression model to describe the relationship. (6 Marks)
- iii) Predict the final score (**Y**) for a patient whose preliminary score (**X**) is **65**.

(2 Marks)

- b) The following are the numbers of patients seen in the outpatient section of a clinic for 10 consecutive working days; 11, 12, 15, 13, 16, 23, 18, 31, 88, 21:

Use the data to answer the following questions:

- i) Find the mean and the median of the data and state with reasons which one of the two is a more appropriate measure of center for this type of data. (2 Marks)

Determine the following measures of dispersion:

- ii) Standard deviation. (2 marks)  
iii) Coefficient of variation. (2 Marks)  
iv) The Inter-quartile range. (2 Marks)
- c) Represent the data in a Box and whisker diagram. (2 Marks)

14.

- a) In 2024, Kitale County Referral Hospital had allocated 92 beds and 27 cots. During the same year, the hospital admitted 3,144 patients and recorded 3,242 discharges and deaths. The total number of inpatient days for the year was 28,600.

Using this information, calculate the following for administrative planning and evaluation at Kitale County Referral Hospital.

- i) Available bed days. (2 Marks)  
ii) Excess or vacant bed days. (2 Marks)  
iii) Daily bed occupancy rate. (2 Marks)  
iv) Turnover Interval. (2 Marks)  
v) Turnover Per Bed. (2 Marks)

- b) Describe the steps followed when performing hypothesis testing in the analysis of health-related data. (3 Marks)
- c) A study was conducted to investigate the effectiveness of hypnotism in reducing pain. Results for randomly selected subjects is shown in the table below. A lower score indicates less pain. The before value is matched to an after value and the differences are calculated. Are the sensory measurements, on average, lower after hypnotism? Test at 5% significance level. (7 Marks)

Subject	A	B	C	D	E	F	G	H
Before	6.6	6.5	9.0	10.3	11.3	8.1	6.3	11.6
After	6.8	2.4	7.4	8.5	8.1	6.1	3.4	2.0

15.

- a) In the year 1990, the surgical department of a county referral hospital recorded various statistics related to inpatient care services. The department was originally allocated 56 beds for the year. However, due to renovations, 20 of these beds were closed between 1st June and 13th August, inclusive. Throughout the year, the department recorded a total of 17,062 occupied bed-days. The total number of patients discharged during the year was 1,921. As a community health officer reviewing this hospital data for planning and reporting purposes, you are required to:
- Calculate the average number of occupied beds during the year. (2 Marks)
  - Calculate the average number of available beds after accounting for the temporary bed closures. (2 Marks)
  - Determine the percentage bed occupancy rate for the year. (2 Marks)
  - Average duration of stay. (2 Marks)
  - Turnover interval. (2 Marks)
  - Turnover per bed. (2 Marks)

- b) Health facilities regularly collect and compile data for use in statistical reporting. As a student of Biostatistics, you are required to understand how this data is transformed into meaningful health information. Describe how you would compile a health information statistical report in a community health setting. Your response should include the process of data collection, organization, analysis, and presentation, as well as the role of the report in improving community health services. (8 Marks)

AMNU

**STANDARD NORMAL DISTRIBUTION: Table Values Represent AREA to the LEFT of the Z score.**

Z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.9	.00005	.00005	.00004	.00004	.00004	.00004	.00004	.00004	.00003	.00003
-3.8	.00007	.00007	.00007	.00006	.00006	.00006	.00006	.00005	.00005	.00005
-3.7	.00011	.00010	.00010	.00010	.00009	.00009	.00008	.00008	.00008	.00008
-3.6	.00016	.00015	.00015	.00014	.00014	.00013	.00013	.00012	.00012	.00011
-3.5	.00023	.00022	.00022	.00021	.00020	.00019	.00019	.00018	.00017	.00017
-3.4	.00034	.00032	.00031	.00030	.00029	.00028	.00027	.00026	.00025	.00024
-3.3	.00048	.00047	.00045	.00043	.00042	.00040	.00039	.00038	.00036	.00035
-3.2	.00069	.00066	.00064	.00062	.00060	.00058	.00056	.00054	.00052	.00050
-3.1	.00097	.00094	.00090	.00087	.00084	.00082	.00079	.00076	.00074	.00071
-3.0	.00135	.00131	.00126	.00122	.00118	.00114	.00111	.00107	.00104	.00100
-2.9	.00187	.00181	.00175	.00169	.00164	.00159	.00154	.00149	.00144	.00139
-2.8	.00256	.00248	.00240	.00233	.00226	.00219	.00212	.00205	.00199	.00193
-2.7	.00347	.00336	.00326	.00317	.00307	.00298	.00289	.00280	.00272	.00264
-2.6	.00466	.00453	.00440	.00427	.00415	.00402	.00391	.00379	.00368	.00357
-2.5	.00621	.00604	.00587	.00570	.00554	.00539	.00523	.00508	.00494	.00480
-2.4	.00820	.00798	.00776	.00755	.00734	.00714	.00695	.00676	.00657	.00639
-2.3	.01072	.01044	.01017	.00990	.00964	.00939	.00914	.00889	.00866	.00842
-2.2	.01390	.01355	.01321	.01287	.01255	.01222	.01191	.01160	.01130	.01101
-2.1	.01786	.01743	.01700	.01659	.01618	.01578	.01539	.01500	.01463	.01426
-2.0	.02275	.02222	.02169	.02118	.02068	.02018	.01970	.01923	.01876	.01831
-1.9	.02872	.02807	.02743	.02680	.02619	.02559	.02500	.02442	.02385	.02330
-1.8	.03593	.03515	.03438	.03362	.03288	.03216	.03144	.03074	.03005	.02938
-1.7	.04457	.04363	.04272	.04182	.04093	.04006	.03920	.03836	.03754	.03673
-1.6	.05480	.05370	.05262	.05155	.05050	.04947	.04846	.04746	.04648	.04551
-1.5	.06681	.06552	.06426	.06301	.06178	.06057	.05938	.05821	.05705	.05592
-1.4	.08076	.07927	.07780	.07636	.07493	.07353	.07215	.07078	.06944	.06811
-1.3	.09680	.09510	.09342	.09176	.09012	.08851	.08691	.08534	.08379	.08226
-1.2	.11507	.11314	.11123	.10935	.10749	.10565	.10383	.10204	.10027	.09853
-1.1	.13567	.13350	.13136	.12924	.12714	.12507	.12302	.12100	.11900	.11702
-1.0	.15866	.15625	.15386	.15151	.14917	.14686	.14457	.14231	.14007	.13786
-0.9	.18406	.18141	.17879	.17619	.17361	.17106	.16853	.16602	.16354	.16109
-0.8	.21186	.20897	.20611	.20327	.20045	.19766	.19489	.19215	.18943	.18673
-0.7	.24196	.23885	.23576	.23270	.22965	.22663	.22363	.22065	.21770	.21476
-0.6	.27425	.27093	.26763	.26435	.26109	.25785	.25463	.25143	.24825	.24510
-0.5	.30854	.30503	.30153	.29806	.29460	.29116	.28774	.28434	.28096	.27760
-0.4	.34458	.34090	.33724	.33360	.32997	.32636	.32276	.31918	.31561	.31207
-0.3	.38209	.37828	.37448	.37070	.36693	.36317	.35942	.35569	.35197	.34827
-0.2	.42074	.41683	.41294	.40905	.40517	.40129	.39743	.39358	.38974	.38591
-0.1	.46017	.45620	.45224	.44828	.44433	.44038	.43644	.43251	.42858	.42465
-0.0	.50000	.49601	.49202	.48803	.48405	.48006	.47608	.47210	.46812	.46414

**STANDARD NORMAL DISTRIBUTION: Table Values Represent AREA to the LEFT of the Z score.**


Z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.50000	.50399	.50798	.51197	.51595	.51994	.52392	.52790	.53188	.53586
0.1	.53983	.54380	.54776	.55172	.55567	.55962	.56356	.56749	.57142	.57535
0.2	.57926	.58317	.58706	.59095	.59483	.59871	.60257	.60642	.61026	.61409
0.3	.61791	.62172	.62552	.62930	.63307	.63683	.64058	.64431	.64803	.65173
0.4	.65542	.65910	.66276	.66640	.67003	.67364	.67724	.68082	.68439	.68793
0.5	.69146	.69497	.69847	.70194	.70540	.70884	.71226	.71566	.71904	.72240
0.6	.72575	.72907	.73237	.73565	.73891	.74215	.74537	.74857	.75175	.75490
0.7	.75804	.76115	.76424	.76730	.77035	.77337	.77637	.77935	.78230	.78524
0.8	.78814	.79103	.79389	.79673	.79955	.80234	.80511	.80785	.81057	.81327
0.9	.81594	.81859	.82121	.82381	.82639	.82894	.83147	.83398	.83646	.83891
1.0	.84134	.84375	.84614	.84849	.85083	.85314	.85543	.85769	.85993	.86214
1.1	.86433	.86650	.86864	.87076	.87286	.87493	.87698	.87900	.88100	.88298
1.2	.88493	.88686	.88877	.89065	.89251	.89435	.89617	.89796	.89973	.90147
1.3	.90320	.90490	.90658	.90824	.90988	.91149	.91309	.91466	.91621	.91774
1.4	.91924	.92073	.92220	.92364	.92507	.92647	.92785	.92922	.93056	.93189
1.5	.93319	.93448	.93574	.93699	.93822	.93943	.94062	.94179	.94295	.94408
1.6	.94520	.94630	.94738	.94845	.94950	.95053	.95154	.95254	.95352	.95449
1.7	.95543	.95637	.95728	.95818	.95907	.95994	.96080	.96164	.96246	.96327
1.8	.96407	.96485	.96562	.96638	.96712	.96784	.96856	.96926	.96995	.97062
1.9	.97128	.97193	.97257	.97320	.97381	.97441	.97500	.97558	.97615	.97670
2.0	.97725	.97778	.97831	.97882	.97932	.97982	.98030	.98077	.98124	.98169
2.1	.98214	.98257	.98300	.98341	.98382	.98422	.98461	.98500	.98537	.98574
2.2	.98610	.98645	.98679	.98713	.98745	.98778	.98809	.98840	.98870	.98899
2.3	.98928	.98956	.98983	.99010	.99036	.99061	.99086	.99111	.99134	.99158
2.4	.99180	.99202	.99224	.99245	.99266	.99286	.99305	.99324	.99343	.99361
2.5	.99379	.99396	.99413	.99430	.99446	.99461	.99477	.99492	.99506	.99520
2.6	.99534	.99547	.99560	.99573	.99585	.99598	.99609	.99621	.99632	.99643
2.7	.99653	.99664	.99674	.99683	.99693	.99702	.99711	.99720	.99728	.99736
2.8	.99744	.99752	.99760	.99767	.99774	.99781	.99788	.99795	.99801	.99807
2.9	.99813	.99819	.99825	.99831	.99836	.99841	.99846	.99851	.99856	.99861
3.0	.99865	.99869	.99874	.99878	.99882	.99886	.99889	.99893	.99896	.99900
3.1	.99903	.99906	.99910	.99913	.99916	.99918	.99921	.99924	.99926	.99929
3.2	.99931	.99934	.99936	.99938	.99940	.99942	.99944	.99946	.99948	.99950
3.3	.99952	.99953	.99955	.99957	.99958	.99960	.99961	.99962	.99964	.99965
3.4	.99966	.99968	.99969	.99970	.99971	.99972	.99973	.99974	.99975	.99976
3.5	.99977	.99978	.99978	.99979	.99980	.99981	.99981	.99982	.99983	.99983
3.6	.99984	.99985	.99985	.99986	.99986	.99987	.99987	.99988	.99988	.99989
3.7	.99989	.99990	.99990	.99990	.99991	.99991	.99992	.99992	.99992	.99992
3.8	.99993	.99993	.99993	.99994	.99994	.99994	.99994	.99995	.99995	.99995
3.9	.99995	.99995	.99996	.99996	.99996	.99996	.99996	.99996	.99997	.99997

**t Table**

cum. prob	$t_{.50}$	$t_{.75}$	$t_{.80}$	$t_{.85}$	$t_{.90}$	$t_{.95}$	$t_{.975}$	$t_{.99}$	$t_{.995}$	$t_{.999}$	$t_{.9995}$
one-tail	<b>0.50</b>	<b>0.25</b>	<b>0.20</b>	<b>0.15</b>	<b>0.10</b>	<b>0.05</b>	<b>0.025</b>	<b>0.01</b>	<b>0.005</b>	<b>0.001</b>	<b>0.0005</b>
two-tails	<b>1.00</b>	<b>0.50</b>	<b>0.40</b>	<b>0.30</b>	<b>0.20</b>	<b>0.10</b>	<b>0.05</b>	<b>0.02</b>	<b>0.01</b>	<b>0.002</b>	<b>0.001</b>
df											
1	0.000	1.000	1.376	1.963	3.078	6.314	12.71	31.82	63.66	318.31	636.62
2	0.000	0.816	1.061	1.386	1.886	2.920	4.303	6.965	9.925	22.327	31.599
3	0.000	0.765	0.978	1.250	1.638	2.353	3.182	4.541	5.841	10.215	12.924
4	0.000	0.741	0.941	1.190	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	0.000	0.727	0.920	1.156	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	0.000	0.718	0.906	1.134	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	0.000	0.711	0.896	1.119	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	0.000	0.706	0.889	1.108	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	0.000	0.703	0.883	1.100	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	0.000	0.700	0.879	1.093	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	0.000	0.697	0.876	1.088	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	0.000	0.695	0.873	1.083	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	0.000	0.694	0.870	1.079	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	0.000	0.692	0.868	1.076	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	0.000	0.691	0.866	1.074	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	0.000	0.690	0.865	1.071	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	0.000	0.689	0.863	1.069	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	0.000	0.688	0.862	1.067	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	0.000	0.688	0.861	1.066	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	0.000	0.687	0.860	1.064	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	0.000	0.686	0.859	1.063	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	0.000	0.686	0.858	1.061	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	0.000	0.685	0.858	1.060	1.319	1.714	2.069	2.500	2.807	3.485	3.768
24	0.000	0.685	0.857	1.059	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	0.000	0.684	0.856	1.058	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	0.000	0.684	0.856	1.058	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	0.000	0.684	0.855	1.057	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	0.000	0.683	0.855	1.056	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	0.000	0.683	0.854	1.055	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	0.000	0.683	0.854	1.055	1.310	1.697	2.042	2.457	2.750	3.385	3.646
40	0.000	0.681	0.851	1.050	1.303	1.684	2.021	2.423	2.704	3.307	3.551
60	0.000	0.679	0.848	1.045	1.296	1.671	2.000	2.390	2.660	3.232	3.460
80	0.000	0.678	0.846	1.043	1.292	1.664	1.990	2.374	2.639	3.195	3.416
100	0.000	0.677	0.845	1.042	1.290	1.660	1.984	2.364	2.626	3.174	3.390
1000	0.000	0.675	0.842	1.037	1.282	1.646	1.962	2.330	2.581	3.098	3.300
<b>Z</b>	0.000	0.674	0.842	1.036	1.282	1.645	1.960	2.326	2.576	3.090	3.291
	0%	50%	60%	70%	80%	90%	95%	98%	99%	99.8%	99.9%
	<b>Confidence Level</b>										



### Chi-Square Distribution Table



The shaded area is equal to  $\alpha$  for  $\chi^2 = \chi^2_{\alpha}$ .

df	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$	$\chi^2_{.100}$	$\chi^2_{.050}$	$\chi^2_{.025}$	$\chi^2_{.010}$	$\chi^2_{.005}$
1	0.000	0.000	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.070	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.300
13	3.565	4.107	5.009	5.892	7.042	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.408	7.564	8.672	10.085	24.769	27.587	30.191	33.409	35.718
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.844	7.633	8.907	10.117	11.651	27.204	30.144	32.852	36.191	38.582
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.034	8.897	10.283	11.591	13.240	29.615	32.671	35.479	38.932	41.401
22	8.643	9.542	10.982	12.338	14.041	30.813	33.924	36.781	40.289	42.796
23	9.260	10.196	11.689	13.091	14.848	32.007	35.172	38.076	41.638	44.181
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.559
25	10.520	11.524	13.120	14.611	16.473	34.382	37.652	40.646	44.314	46.928
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.808	12.879	14.573	16.151	18.114	36.741	40.113	43.195	46.963	49.645
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.993
29	13.121	14.256	16.047	17.708	19.768	39.087	42.557	45.722	49.588	52.336
30	13.787	14.953	16.791	18.493	20.599	40.256	43.773	46.979	50.892	53.672
40	20.707	22.164	24.433	26.509	29.051	51.805	55.758	59.342	63.691	66.766
50	27.991	29.707	32.357	34.764	37.689	63.167	67.505	71.420	76.154	79.490
60	35.534	37.485	40.482	43.188	46.459	74.397	79.082	83.298	88.379	91.952
70	43.275	45.442	48.758	51.739	55.329	85.527	90.531	95.023	100.425	104.215
80	51.172	53.540	57.153	60.391	64.278	96.578	101.879	106.629	112.329	116.321
90	59.196	61.754	65.647	69.126	73.291	107.565	113.145	118.136	124.116	128.299
100	67.328	70.065	74.222	77.929	82.358	118.498	124.342	129.561	135.807	140.169